

Supplementary Table 2. Clinical outcomes according to timing of enteral nutrition in patients without shockable initial rhythm

Variable	Delayed nutrition support (n=19)	Early nutrition support (n=16)	Total (n=35)	P-value
Intensive care (day)	6.7±5.3	11.9±7.8	9.2±7.0	0.023
Total hospital stay (day) ^a	6.0±5.6	17.0±46.0	11.5±33.0	0.088
APACHE II score	21.7±9.9	16.8±6.9	19.4±8.9	0.097
Survival to discharge	7 (36.8)	14 (82.4)	21 (58.3)	<0.006
Good neurologic outcome ^b	2 (10.5)	8 (47.1)	10 (27.8)	0.025
Complication				
Aspiration pneumonia	4 (21.1)	4 (23.5)	8 (22.2)	1
Regurgitation	2 (10.5)	2 (11.8)	4 (11.1)	1
Ileus	1 (5.3)	0	1 (2.8)	1
Fever	10 (52.6)	5 (29.4)	15 (41.7)	0.158
Hypoglycemia	4 (21.1)	4 (23.5)	8 (22.2)	1
Hyperglycemia	7 (36.8)	11 (64.7)	18 (50.0)	0.095

Values are presented as mean±standard deviation or number (%).

APACHE: Acute physiology and chronic health evaluation.

a) Median±standard deviation; b) Cerebral performance category ≤2.