

Supplementary Table 7. Clinical parameters by group during the study period

Parameter	Group 1 (n=41)	Group 2 (n=54)	P-value
Day 4			
Mean arterial pressure (mm Hg)	69 (64–80) (n=34)	69 (60–80) (n=53)	0.18
PaO ₂ /FiO ₂	220 (170–286) (n=30)	228 (193–315) (n=49)	0.28
Creatinine (mg/dl)	1.1 (0.7–1.4) (n=34)	0.8 (0.6–1.3) (n=53)	0.18
Lymphocyte count (/mm ³)	643±48	751±708	0.63
Total bilirubin (mg/dl)	0.6 (0.4–1.0) (n=34)	0.7 (0.5–1.5) (n=53)	0.19
C-reactive protein (mg/L)	123±80	150±99	0.4
Lactate (mmol/L)	1.8 (1.3–3.7)	1.7 (1.4–2.5)	0.59
Norepinephrine equivalent dose (µg/kg/min)	0.10±0.21	0.06±0.23	0.03
SOFA score	11±7	8±4	0.05
Net fluid retention (ml) ^{a)}	180 (–338 to 1,053) (n=31)	315 (–392 to 785) (n=54)	0.62
Day 7			
Mean arterial pressure (mm Hg)	68 (57–78) (n=32)	72 (63–80) (n=53)	0.18
PaO ₂ /FiO ₂	237 (154–285) (n=23)	236 (180–318) (n=41)	0.38
Creatinine (mg/dl)	0.8 (0.5–1.2) (n=32)	0.7 (0.5–1.1) (n=53)	0.72
Lymphocyte count (/mm ³)	792±747	1,082±1,091	0.11
Total bilirubin (mg/dl)	0.8 (0.5–1.3) (n=32)	0.8 (0.4–1.4) (n=53)	0.94
C-reactive protein (mg/L)	103±89	54±50	0.02
Lactate (mmol/L)	1.6 (1.1–2.4)	1.7 (1.2–2.5)	0.62
Norepinephrine equivalent dose (µg/kg/min)	0.10±0.22	0.05±0.22	0.27
SOFA score	11±7	8±4	0.01
Net fluid retention (ml) ^{a)}	279 (–341 to 725) (n=28)	160 (–769 to 850) (n=49)	0.73
Change relative to day 1			
Lymphocyte count at day 7 (/mm ³)	102±754	272±1,517	0.29
C-reactive protein at day 7 (mg/L)	–98±126	–134±123	0.13

Values are presented as median (interquartile range) or mean±standard deviation. Group 1: ____; Group 2: ____.

PaO₂: arterial partial pressure of oxygen; FiO₂: fraction of inspired oxygen; SOFA: Sequential Organ Failure Assessment.

a) Net fluid retention is calculated as the difference between intake and output of all fluids (urine volume, dialysis volume, drainage volume, and stool weight).